

Practical Statistics Problems, Imperial College, Nov 2008

- 1) An experiment is searching for quarks of charge $2/3$, which are expected to produce $4/9$ the ionisation I_0 of unit charged particles. In an exposure in which 10^5 cosmic ray tracks are observed, 1 track has its ionisation measured as $0.44I_0$. The detector is such that ionisation measurements are Gaussian distributed about their true values with standard deviation σ . Calculate the probability that this could be a statistical fluctuation on the ionisation of a unit charged particle for the following different assumptions:
 - a) $\sigma = 0.07I_0$ for all 10^5 track,
 - b) For 99% of the tracks $\sigma = 0.07I_0$, while for the remainder it is $0.14I_0$.

- 2) An experiment is determining the decay rate λ for a new particle X, whose probability density for decay at time t is proportional to $\exp(-\lambda t)$. A total of nine decays are observed at decay times 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 and 0.9 picoseconds. Calculate the likelihood function $L(\lambda)$ at suitable values of λ (most easily done by a simple computer program), and draw a graph of the results. Find the best estimate of λ from the maximum of the likelihood curve, and a “ $\pm\sigma$ ” range for λ by finding the values of λ where the logarithm to the base e of the likelihood function decreases by 0.5 units from its maximum value.

- 3) i) A tracker has detector 6 elements at $x = -11, -10, -9, +9, +10$ and $+11$ cms, which each measure a track's y -coordinate to an accuracy of ± 1 cm. A straight line $y = a + bx$ is fitted (for example by chi-squared) to the data from the 3 elements at positive x (L1); a second (L2) for the data at negative x ; and a third (L3) to all 6 detector elements. The inverse error matrix for a and b has elements $M_{aa}^{-2} = \sum 1/\sigma_i^2$ $M_{bb}^{-2} = \sum x_i^2/\sigma_i^2$ $M_{ab}^{-2} = \sum x_i/\sigma_i^2$, where the measurements are $y_i \pm \sigma_i$ at x_i . Evaluate the error matrix for a and b for each of the 3 fits. How do the errors and correlations compare with what you expect?
 - ii) When two measurements for a pair of quantities are combined optimally, the errors on the combined parameters are such that $M_c = M_1 + M_2$, where M_c is the inverse error matrix for the combination, and M_1 and M_2 are those for the separate measurements. Determine the error matrix for the combination of the parameters of L1 and L2. Explain why the errors for the combination are considerably smaller than those for L1 and L2 separately.

- 4) The coverage $C(\mu)$ is a property of a statistical technique for estimating a range for a parameter μ at a confidence level α (e.g. 68%, 90% or whatever). It is the fraction of times that, in repetitions of the procedure with different data, the estimated range contains the true value μ . In a Poisson counting experiment with n observed events, one method of estimating a range for the Poisson parameter μ uses the estimate $n \pm \sqrt{n}$ i.e. from $n - \sqrt{n}$ to $n + \sqrt{n}$. This is

supposed to have 68% coverage. Determine the actual coverage $C(\mu)$ at $\mu = 3.41$ and 3.42 as follows:

Determine for which measured values, the nominal range from the " $n \pm \sqrt{n}$ " procedure includes the specified true value, and then add up the Poisson probabilities for obtaining these measured values, again assuming the specified value of the Poisson parameter.

Explain why plots of the coverage $C(\mu)$ as a function of the Poisson parameter value μ have discontinuities.

- 5) An experiment is searching for the SM Higgs. With no Higgs production, 100 events are expected; if the Higgs is produced, 110 events are expected. The experiment observes 130 events, which is 3σ above the 'No Higgs' prediction, so the p-value for the null hypothesis is 0.1%. The Lab Publicity Officer announces that we now are 99.9% certain that the Higgs has been discovered.

Comment.